



## Arctic marine gravity and bathymetry from 3 years of Cryosat-2 SAR altimetry (DTU13 Gravity)

Andersen, Ole Baltazar; Stenseng, Lars; Knudsen, Per; Jain, Maulik

*Publication date:*  
2013

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Andersen, O. B., Stenseng, L., Knudsen, P., & Jain, M. (2013). *Arctic marine gravity and bathymetry from 3 years of Cryosat-2 SAR altimetry (DTU13 Gravity)*. Abstract from AGU Fall Meeting 2013, San Francisco, United States. <http://abstractsearch.agu.org/meetings/2013/FM/sections/G/sessions/G11B/abstracts/G11B-0922.html>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**CONTROL ID:** 1817993

**TITLE:** Arctic marine gravity and bathymetry from 3 years of Cryosat-2 SAR altimetry (DTU13 Gravity)

**AUTHORS (FIRST NAME, LAST NAME):** Ole Baltazar Andersen<sup>1</sup>, Lars Stenseng<sup>1</sup>, Per Knudsen<sup>1</sup>, Maulik Jain<sup>1</sup>

**INSTITUTIONS (ALL):** 1. DTU Space, Lyngby, Denmark.

**ABSTRACT BODY:** The accuracy of the Arctic marine gravity field has for many been severely limited by the availability and accuracy of altimeter data in the Arctic Ocean. Until recently only ERS-1 provided non-repeat (0.9 year) geodetic mission altimetry in the Arctic Ocean and only up to 82N.

With the launch of Cryosat-2 three years ago a new source of high quality altimetric data has become available. The Cryosat-2 delay Doppler altimeter offers a factor of 20 improvements in along track resolution, an along-track footprint length that does not vary with wave height (sea state) and at least a factor of two in sea surface height precision. Over the Arctic Ocean the Cryosat-2 generally operates in SAR altimetry mode for cryospheric studies. We have tested the standard ESA L2 SAR altimetric data for the first 3 years and developed robust empirical retracers for ice-covered regions and processing 3 years of L1 SAR altimetry in the Arctic Ocean for gravity field determination. Extensive testing, interpretation and improvement of methods to handles the new class of data has been investigated and the first result from a new Arctic Ocean wide gravity field will be presented as well as initial test of derived altimetric bathymetry using the new gravity field data.

**KEYWORDS:** 1219 GEODESY AND GRAVITY Gravity anomalies and Earth structure, 1241 GEODESY AND GRAVITY Satellite geodesy: technical issues, 1640 GLOBAL CHANGE Remote sensing.

(No Image Selected)

(No Table Selected)

### **Additional Details**

**Previously Presented Material:**

### **Contact Details**

**CONTACT (NAME ONLY):** Ole Andersen

**CONTACT (E-MAIL ONLY):** oa@space.dtu.dk

**TITLE OF TEAM:**

---